

A NEW MILKY WAY DWARF SATELLITE IN CANES VENATICI

D. B. ZUCKER,¹ V. BELOKUROV,¹ N. W. EVANS,¹ M. I. WILKINSON,¹ M. J. IRWIN,¹ T. SIVARANI,² S. HODGKIN,¹
D. M. BRAMICH,¹ J. M. IRWIN,¹ G. GILMORE,¹ B. WILLMAN,³ S. VIDRIH,¹ M. FELLHAUER,¹ P. C. HEWETT,¹ T. C. BEERS,²
E. F. BELL,⁴ E. K. GREBEL,⁵ D. P. SCHNEIDER,⁶ H. J. NEWBERG,⁷ R. F. G. WYSE,⁸ C. M. ROCKOSI,⁹ B. YANNY,¹⁰ R. LUPTON,¹¹
J. A. SMITH,¹² J. C. BARENTINE,¹³ H. BREWINGTON,¹³ J. BRINKMANN,¹³ M. HARVANEK,¹³ S. J. KLEINMAN,¹³ J. KRZESINSKI,^{13,14}
D. LONG,¹³ A. NITTA,¹³ AND S. A. SNEDDEN¹³

Received 2006 April 10; accepted 2006 April 19; published 2006 May 18

ABSTRACT

In this Letter, we announce the discovery of a new dwarf satellite of the Milky Way, located in the constellation Canes Venatici. It was found as a stellar overdensity in the north Galactic cap using Sloan Digital Sky Survey Data Release 5 (SDSS DR5). The satellite’s color-magnitude diagram shows a well-defined red giant branch as well as a horizontal branch. As judged from the tip of the red giant branch, it lies at a distance of ~ 220 kpc. Based on the SDSS data, we estimate an absolute magnitude of $M_V \sim -7.9$, a central surface brightness of $\mu_{0,V} \sim 28$ mag arcsec⁻², and a half-light radius of ~ 8.5 (~ 550 pc at the measured distance). The outer regions of Canes Venatici appear extended and distorted. The discovery of such a faint galaxy in proximity to the Milky Way strongly suggests that more such objects remain to be found.

Subject headings: galaxies: dwarf — galaxies: individual (Canes Venatici) — Local Group

1. INTRODUCTION

There are 10 known dwarf spheroidal (dSph) companions of the Milky Way. Together with the two dwarf irregulars (the Large and Small Magellanic Clouds), these make up all the known satellite galaxies of the Milky Way. The dSphs have such low surface brightness that they have often been found serendipitously. For example, while Sextans (Irwin et al. 1990) was found as part of an automated search, the intrinsically far more luminous Sagittarius dSph was first identified kinematically from a radial velocity survey of stars in Galactic center fields (Ibata et al. 1995).

As such galaxies are resolvable into individual stars because of their proximity, they are detectable as enhancements in the stellar number density in large photometric surveys. For example, Martin et al. (2004) analyzed overdensities of M giants in the Two Micron All Sky Survey (2MASS) and claimed the detection of a new disrupted dSph in Canis Major, although this remains controversial. Willman et al. (2005) systematically

surveyed ~ 5800 deg² of the Sloan Digital Sky Survey (SDSS; York et al. 2000) and identified a dSph companion to the Milky Way in the constellation of Ursa Major. Subsequent spectroscopic observations (Kleyna et al. 2005) showed that Ursa Major was a dark matter-dominated dSph. Meanwhile, Zucker et al. (2004, 2006) announced the discovery of Andromeda IX and X, new dSph satellites of M31 found in SDSS data with similar methods.

Very recently, Belokurov et al. (2006b) mapped out the stars satisfying the color cut $g - r < 0.4$ in almost all of SDSS Data Release 5 (DR5). They dubbed this plot of the high-latitude Galactic northern hemisphere the “Field of Streams,” owing to its wealth of prominent stellar substructure. Visible by eye in the plot is a heretofore undetected enhancement of the stellar density, located in the constellation Canes Venatici. Here, we show that this corresponds to a new dSph companion—the eleventh—of the Milky Way.

2. DATA AND DISCOVERY

The SDSS is an imaging and spectroscopic survey. SDSS imaging data are produced in five photometric bands, namely, u , g , r , i , and z (Fukugita et al. 1996; Gunn et al. 1998; Hogg et al. 2001; Adelman-McCarthy et al. 2006; Gunn et al. 2006), and are automatically processed through pipelines to measure photometric and astrometric properties (Lupton et al. 1999; Stoughton et al. 2002; Smith et al. 2002; Pier et al. 2003; Ivezić et al. 2004). DR5 primarily covers ~ 8000 deg² around the north Galactic pole (NGP).

In the process of analyzing this area of the NGP (Belokurov et al. 2006b), we visually identified a stellar overdensity in the constellation Canes Venatici. The upper left panel of Figure 1 shows a gray-scale SDSS image of the sky centered on the stellar overdensity; note that no obvious object can be seen. However, a roughly elliptical overdensity of objects classified by the SDSS pipeline as stars is readily visible in the photometric data (*upper middle and right panels*). Plotting these stars in a color-magnitude diagram (CMD) reveals a clear red giant branch (RGB) and horizontal branch (*lower panels*). This CMD appears similar to that seen in SDSS data for the Sextans dSph

¹ Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK; zucker@ast.cam.ac.uk, vasily@ast.cam.ac.uk, nwe@ast.cam.ac.uk.

² Department of Physics and Astronomy, Center for the Study of Cosmic Evolution, and Joint Institute for Nuclear Astrophysics, Michigan State University, East Lansing, MI 48824.

³ Center for Cosmology and Particle Physics, Department of Physics, New York University, 4 Washington Place, New York, NY 10003.

⁴ Max Planck Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany.

⁵ Astronomical Institute of the University of Basel, Department of Physics and Astronomy, Venusstrasse 7, CH-4102 Binningen, Switzerland.

⁶ Department of Astronomy and Astrophysics, Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802.

⁷ Rensselaer Polytechnic Institute, Troy, NY 12180.

⁸ Johns Hopkins University, 3701 San Martin Drive, Baltimore, MD 21218.

⁹ Lick Observatory, University of California, Santa Cruz, CA 95064.

¹⁰ Fermi National Accelerator Laboratory, P.O. Box 500, Batavia, IL 60510.

¹¹ Princeton University Observatory, Peyton Hall, Princeton, NJ 08544.

¹² Los Alamos National Laboratory, ISR-4, MS D448, Los Alamos, NM 87545.

¹³ Apache Point Observatory, P.O. Box 59, Sunspot, NM 88349.

¹⁴ Mount Suhora Observatory, Cracow Pedagogical University, ul. Podchorążych 2, 30-084 Cracow, Poland.




FIG. 1.—Canes Venatici dwarf. *Upper left*: Combined SDSS g, r, i images of a $1^\circ \times 1^\circ$ field centered on the overdensity. $\Delta\alpha$ and $\Delta\delta$ are the relative offsets in right ascension and declination, measured in degrees. The dotted circle indicates a radius of 0.25° . *Upper middle*: Spatial distribution of all objects classified as stars in the same area. *Upper right*: Binned spatial density of all stellar objects. The inner dotted circle marks a radius of 0.25° , the middle circle a radius of 0.5° , and the outermost circle a radius of 0.56° . Bins are $0.05^\circ \times 0.05^\circ$, smoothed with a Gaussian with an FWHM of 0.1° . *Lower left*: CMD of all stellar objects within the inner 0.25° radius circle; note the clear RGB, with hints of a horizontal branch, even without removal of field contamination. The filled star shows the location of a carbon star in the field of CVn (see text). *Lower middle*: CMD of a comparably sized control area, showing all stellar objects between 0.5° (middle circle) and 0.56° (outer circle) from the center. *Lower right*: Color-magnitude density plot (Hess diagram), showing the inner CMD minus a control CMD taken between 0.5° and 2° from the center, normalized to the number of stars in each CMD. RGB and horizontal-branch fiducial ridgelines derived from SDSS photometry of the globular cluster NGC 2419 ($[\text{Fe}/\text{H}] \sim -2.1$) are overplotted as dashed white-and-black lines; RGB fiducials for M3 ($[\text{Fe}/\text{H}] \sim -1.6$) and M71 ($[\text{Fe}/\text{H}] \sim -0.7$) from Clem (2005) are shown in dotted white-and-black and solid black lines, respectively. All fiducials were corrected for Galactic foreground extinction (Schlegel et al. 1998) and shifted to a distance modulus of 21.75. The thin dashed vertical line shows the cut $g - i < -0.1$ used to select candidate blue horizontal-branch stars.

(see, e.g., Fig. 1 of Willman et al. 2005), although considerably more distant. Based on its apparent morphology and on the presence of a distinct stellar population, we conclude that this is most likely a hitherto unknown dSph galaxy. As is customary, we name it after its constellation, Canes Venatici (CVn).

3. PHYSICAL PROPERTIES AND STELLAR POPULATION

The distance of a distinct, metal-poor stellar population can be estimated from the I -band magnitude of the tip of the red giant branch (TRGB; e.g., Mould et al. 1983; Lee et al. 1993a). We applied the transforms from Smith et al. (2002) to convert dereddened SDSS g, r, i magnitudes to V, I magnitudes. The cumulative I -band luminosity function for the RGB shows a sharp rise starting at $I \sim 17.75$ (which corresponds to $i \sim 18.23$). The horizontal-branch fit in the lower right panel of Figure 1 provides supporting evidence for the reliability of our procedure. Assuming a metallicity of $[\text{Fe}/\text{H}] \sim -2$ (see below) and a TRGB color of $(V - I)_{\text{TRGB}} = 1.4$, from the calibration of Da Costa & Armandroff (1990) we obtain the relation $(m - M)_0 = I_{\text{TRGB}} + 3.97$, yielding a distance modulus of $(m - M)_0 \sim 21.75 \pm 0.20$ ($\sim 220_{-16}^{+25}$ kpc). This is comparable to the most distant Milky Way dSph satellites previously known, Leo I and II (250 and 215 kpc, respectively; Lee et al. 1993b; Lee 1995).

The age and metallicity of the stars in CVn can be estimated by comparison with the CMDs of single-epoch, single-metallicity populations, e.g., Galactic globular clusters. In the lower

right panel of Figure 1, we show a field-star-subtracted Hess diagram (color-magnitude density plot) in $(i, g - i)$. Fiducial ridgelines of three globular clusters—NGC 2419 ($[\text{Fe}/\text{H}] \sim -2.1$), M3 ($[\text{Fe}/\text{H}] \sim -1.6$), and M71 ($[\text{Fe}/\text{H}] \sim -0.7$)—are overplotted from left to right. The best match is to NGC 2419, and in combination with the prominent horizontal branch, the similarity indicates that CVn is dominated by an old, metal-poor ($[\text{Fe}/\text{H}] \sim -2$) stellar population. The horizontal branch extends very far to the blue, although it is predominantly red. Such red-dominated horizontal branches, hinting at a range of ages in the stellar population, are also seen in other metal-poor dSphs (e.g., Harbeck et al. 2001).

To investigate the spatial morphology, we select candidate members by applying a mask built as follows. First, we find the class conditional probabilities P_{CVn} and P_{bg} (see, e.g., Belokurov et al. 2006a) of a star belonging to CVn and the background, respectively. These are just the normalized Hess diagrams. The ratio of the class conditional probabilities $x = P_{\text{CVn}}/P_{\text{bg}}$ is converted to a probability of membership of CVn via $p = x/(1 + x)$. A cut of $p > 0.8$ is used to select stars plotted in the top panel of Figure 2. These objects are binned into 30×30 bins, each $0.033^\circ \times 0.033^\circ$, and smoothed with a Gaussian with FWHM of 0.067° to yield the plot in the lower panel. The density contours, representing 2, 3, 5, 7, 10, 15, 20, and 25 σ above the background level, are markedly irregular, particularly in the outer parts—somewhat reminiscent of the Ursa Minor dSph. The black crosses are candidate blue horizontal-




FIG. 2.—Morphology of the CVn dwarf. *Upper panel:* Spatial distribution of stars selected from CMD regions with the highest contrast between CVn and field stellar populations. $\Delta\alpha$ and $\Delta\delta$ are relative offsets (degrees of arc) in right ascension and declination. *Lower panel:* Contour plot of the spatial distribution of these stars; candidate blue horizontal-branch stars (from the area blueward of the dashed line in the lower right panel of Fig. 1) are overplotted with crosses. The contours are 2, 3, 5, 7, 10, 15, 20, and 25 σ above the background level. The location of the carbon star, SDSS J132755.56+333521.7, is indicated by the black star.

branch (BHB) stars selected with $g - i < -0.1$. The distribution of BHB stars is elongated in the same manner as the density contours and traces the same underlying morphology.

To estimate the properties listed in Table 1, we first analyze the morphology shown in Figure 2 to derive the centroid from the density-weighted first moment of the distribution, and the average ellipticity and position angle using the three density-weighted second moments (e.g., Stobie 1980). The radial profile shown in Figure 3 is derived by computing the average density within elliptical annuli after first subtracting a constant asymptotic background level (0.25 arcmin^{-2}) reached at large radii. We then fit the radial profile with standard Plummer and exponential laws (Fig. 3; see also Irwin & Hatzidimitriou 1995). The best-fitting position angle, ellipticity, and half-light radii are listed in Table 1. At the distance of CVn, the approximately 8.5 half-light radius corresponds to $\sim 550 \text{ pc}$. The overall extent is $\sim 2 \text{ kpc}$ along the major axis, making CVn one of the largest of the Milky Way dSphs. Although Plummer and exponential laws provide reasonable fits to the data, there are some discrepant data points in Figure 3 (cf. Sextans' profile in Irwin & Hatzidimitriou 1995).

The overall luminosity is computed by masking the stellar locus of CVn and computing the total flux within the mask and within the elliptical half-light radius. A similar mask, but covering a larger area to minimize shot noise, well outside the main body of CVn, is scaled by relative area and used to compute the foreground contamination within the half-light radius. After correcting for this contamination, the remaining flux is scaled to the total, assuming the fitted profiles are a fair representation of the overall flux distribution. We also apply a correction of 0.5 mag for unresolved/faint stars, based on the stellar luminosity functions of other low-metallicity, low surface brightness dSphs. The resulting luminosity

TABLE 1
PROPERTIES OF THE CANES VENATICI DWARF

Parameter ^a	Value
Coordinates (J2000)	$13^{\text{h}}28^{\text{m}}03^{\text{s}}.5, +33^{\circ}33'21''.0 \pm 10''$
Galactic coordinates (l, b)	$74^{\circ}3, 79^{\circ}8$
Position angle	$73^{\circ} \pm 3^{\circ}$
Ellipticity	0.38
r_h (Plummer)	$8'.5 \pm 0'.5$
r_h (exponential)	$8'.4 \pm 0'.5$
A_V	0.05 mag
$\mu_{0,V}$ (Plummer)	$28.2 \pm 0.5 \text{ mag}$
$\mu_{0,V}$ (exponential)	$27.8 \pm 0.5 \text{ mag}$
V_{tot}	$13.9 \pm 0.5 \text{ mag}$
$(m - M)_0$	$21.75 \pm 0.20 \text{ mag}$
$M_{\text{tot},V}$	$-7.9 \pm 0.5 \text{ mag}$

^a Surface brightnesses and integrated magnitudes are corrected for the mean Galactic foreground reddenings, A_V , shown.

and central surface brightness estimates, $M_{\text{tot},V} \sim -7.9$ and $\mu_{0,V} \sim 28 \text{ mag arcsec}^2$, are roughly consistent with the central surface brightness–absolute magnitude relation noted for Local Group dSph satellites (e.g., Grebel et al. 2003). However, CVn lies far from the central surface brightness–host galaxy distance correlation found for other Local Group dSph satellites (e.g., Fig. 7 of McConnachie & Irwin 2006), suggesting that this correlation may be at least partially due to observational biases (e.g., Willman et al. 2004).

A search of the SDSS DR5 spectroscopic database revealed the spectrum of a CH carbon star, SDSS J132755.56+333521.7, in close proximity to the center of CVn (lower panel of Fig. 2). As shown in Figure 4, the star's spectral features indicate that it is most likely a giant, rather than a dwarf. There are no 2MASS point sources within $15''$, and a comparison of the SDSS coordinates with earlier-epoch APM data is consistent with zero proper motion, both of which would argue in favor of its being a distant object. Finally, the carbon star's dereddened r -band magnitude is 19.18; at the calculated distance modulus for CVn, ~ 21.75 , the star would have an absolute magnitude $M_r \sim -2.5$, fairly typical of a CH carbon giant. Hence, we conclude that SDSS J132755.56+333521.7 is a carbon star in CVn. We measure a heliocentric radial velocity of $+36 \pm 20 \text{ km s}^{-1}$ for the carbon star from cross-correlation with the spectrum of another CH carbon star of known velocity (1249+0146; see Totten & Irwin 1998). The carbon star's radial velocity in the line-of-sight direction (after correction to the Galactic rest frame) is $\sim 80 \pm 20 \text{ km s}^{-1}$. Given




FIG. 3.—Profile of CVn, showing the background-subtracted stellar density in elliptical annuli as a function of mean radius. The left panel is logarithmic in both axes, and the right panel is linear in radius. The overplotted lines are fitted Plummer (solid) and exponential (dashed) profiles.




FIG. 4.—Spectrum of SDSS J132755.56+333521.7 (*bottom*), a carbon star located near the center of CVn (see lower left panel of Fig. 1 and the lower panel of Fig. 2). Spectra of a giant carbon star (SDSS J075116.37+391201.4, *middle*) and a dwarf carbon star (SDSS J001145.30-004710.2, *top*) with similar temperatures ($T_{\text{eff}} \sim 4000$ K; Downes et al. 2004) are shown for comparison. The CN bands are stronger in the reference giant than in the dwarf, while Na I, Cr I, and Mg I *b* lines and the C_2 bands around 6150 \AA are stronger in the dwarf. The strength of the former features and the weakness of the latter features in SDSS J132755.56+333521.7 indicate that it is most likely a giant carbon star in CVn.

the distance of CVn, this is also approximately the radial velocity that would be seen from the Galactic center. We plan to obtain follow-up spectroscopy of RGB stars in CVn to supplement this first estimate of its systemic velocity with a more detailed analysis of the new dwarf's kinematics.

4. CONCLUSIONS

We have discovered a new dwarf spheroidal Milky Way satellite in the constellation Canes Venatici. It has an absolute magnitude of $M_V \sim -7.9$ and a surface brightness of $\mu_{0,V} \sim 28 \text{ mag arcsec}^{-2}$. We find that CVn is dominated by an old, metal-poor ($[\text{Fe}/\text{H}] \sim -2$) stellar population. At a distance of $\sim 220 \text{ kpc}$, CVn is one of the most remote of the known dSph companions to the Milky Way.

It is important to carry out a well-defined census of Milky

Way dwarf galaxies to the faintest possible luminosities. Numerical simulations in cold dark matter cosmological models predict that the Milky Way's halo should contain about 500 satellites comparable to the dSphs (Moore et al. 1999; Klypin et al. 1999). This contrasts with the handful of known Milky Way dSphs. The new dSph in Canes Venatici, taken with the recent discovery of Ursa Major (Willman et al. 2005), emphasizes how incomplete our current knowledge is. Further analysis of photometric data from large-area digital surveys such as SDSS will thus likely lead to the discovery of more dSphs around the Milky Way in the near future.

D. B. Z., V. B., M. I. W., D. M. B., and M. F. acknowledge the financial support of the Particle Physics and Astronomy Research Council of the United Kingdom.

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the US Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England.¹⁵

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, the University of Basel, Cambridge University, Case Western Reserve University, the University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max Planck Institute for Astronomy (MPIA), the Max Planck Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, the University of Pittsburgh, the University of Portsmouth, Princeton University, the US Naval Observatory, and the University of Washington.

¹⁵ The SDSS Web site is <http://www.sdss.org/>.

REFERENCES

- Adelman-McCarthy, J. K., et al. 2006, *ApJS*, 162, 38
 Belokurov, V., Evans, N. W., Irwin, M. J., Hewett, P. C., & Wilkinson, M. I. 2006a, *ApJ*, 637, L29
 Belokurov, V., et al. 2006b, *ApJ*, 642, L137
 Clem, J. L. 2005, Ph.D. thesis, Univ. Victoria
 Da Costa, G. S., & Armandroff, T. E. 1990, *AJ*, 100, 162
 Downes, R. A., et al. 2004, *AJ*, 127, 2838
 Fukugita, M., Ichikawa, T., Gunn, J. E., Doi, M., Shimasaku, K., & Schneider, D. P. 1996, *AJ*, 111, 1748
 Grebel, E. K., Gallagher, J. S., & Harbeck, D. 2003, *AJ*, 125, 1926
 Gunn, J. E., et al. 1998, *AJ*, 116, 3040
 ———. 2006, *AJ*, 131, 2332
 Harbeck, D., et al. 2001, *AJ*, 122, 3092
 Hogg, D. W., Finkbeiner, D. P., Schlegel, D. J., & Gunn, J. E. 2001, *AJ*, 122, 2129
 Ibata, R. A., Gilmore, G., & Irwin, M. J. 1995, *MNRAS*, 277, 781
 Irwin, M. J., Bunclark, P. S., Bridgeland, M. T., & McMahon, R. G. 1990, *MNRAS*, 244, 16P
 Irwin, M., & Hatzidimitriou, D. 1995, *MNRAS*, 277, 1354
 Ivezić, Ž., et al. 2004, *Astron. Nachr.*, 325, 583
 Kleyana, J. T., Wilkinson, M. I., Evans, N. W., & Gilmore, G. 2005, *ApJ*, 630, L141
 Klypin, A., Kravtsov, A. V., Valenzuela, O., & Prada, F. 1999, *ApJ*, 522, 82
 Lee, M. G. 1995, *AJ*, 110, 1155
 Lee, M. G., Freedman, W. L., & Madore, B. F. 1993a, *ApJ*, 417, 553
 Lee, M. G., Freedman, W., Mateo, M., Thompson, I., Roth, M., & Ruiz, M.-T. 1993b, *AJ*, 106, 1420
 Lupton, R., Gunn, J., & Szalay, A. 1999, *AJ*, 118, 1406
 Martin, N. F., Ibata, R. A., Bellazzini, M., Irwin, M. J., Lewis, G. F., & Dehnen, W. 2004, *MNRAS*, 348, 12
 McConnachie, A., & Irwin, M. 2006, *MNRAS*, 365, 1263
 Moore, B., Governato, F., Lake, G., Quinn, T., Stadel, J., & Tozzi, P. 1999, *ApJ*, 524, L19
 Mould, J., Kristian, J., & Da Costa, G. 1983, *ApJ*, 270, 471
 Pier, J. R., Munn, J. A., Hindsley, R. B., Hennessy, G. S., Kent, S. M., Lupton, R. H., & Ivezić, Ž. 2003, *AJ*, 125, 1559
 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
 Smith, J. A., et al. 2002, *AJ*, 123, 2121
 Stobie, R. S. 1980, *J. British Interplanet. Soc.*, 33, 323
 Stoughton, C., et al. 2002, *AJ*, 123, 485
 Totten, E. J., & Irwin, M. J. 1998, *MNRAS*, 294, 1
 Willman, B., Governato, F., Dalcanton, J. J., Reed, D., & Quinn, T. 2004, *MNRAS*, 353, 639
 Willman, B., et al. 2005, *ApJ*, 626, L85
 York, D. G., et al. 2000, *AJ*, 120, 1579
 Zucker, D. B., et al. 2004, *ApJ*, 612, L121
 ———. 2006, *ApJL*, submitted (astro-ph/0601599)